

§ 640.3

(4) The blood is held for observation until a significant inspection consistent with the requirements of § 640.5(e) can be made.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 4015, Jan. 28, 1976; 42 FR 59878, Nov. 22, 1977; 43 FR 34460, Aug. 4, 1978; 49 FR 15187, Apr. 18, 1984; 49 FR 23834, June 8, 1984; 50 FR 4138, Jan. 29, 1985; 53 FR 116, Jan. 5, 1988; 55 FR 11013, Mar. 26, 1990; 63 FR 16685, Apr. 6, 1998; 64 FR 45371, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 31165, June 11, 2001; 66 FR 40889, Aug. 6, 2001; 67 FR 9587, Mar. 4, 2002]

§ 640.3 Suitability of donor.

(a) *Method of determining.* The suitability of a donor as a source of Whole Blood shall be determined by a qualified physician or by persons under his supervision and trained in determining suitability. Such determination shall be made on the day of collection from the donor by means of medical history, a test for hemoglobin level, and such physical examination as appears necessary to a physician who shall be present on the premises when examinations are made, except that the suitability of donors may be determined when a physician is not present on the premises, provided the establishment (1) maintains on the premises, and files with the Center for Biologics Evaluation and Research, a manual of standard procedures and methods, approved by the Director of the Center for Biologics Evaluation and Research, that shall be followed by employees who determine suitability of donors, and (2) maintains records indicating the name and qualifications of the person immediately in charge of the employees who determine the suitability of donors when a physician is not present on the premises.

(b) *Qualifications of donor; general.* Except as provided in paragraph (f) of this section and for autologous donations, a person may not serve as a source of Whole Blood more than once in 8 weeks. In addition, donors shall be in good health, as indicated in part by:

- (1) Normal temperature;
- (2) Demonstration that systolic and diastolic blood pressures are within normal limits, unless the examining physician is satisfied that an individual with blood pressures outside these limits is an otherwise qualified

21 CFR Ch. I (4–1–15 Edition)

donor under the provisions of this section;

(3) For allogeneic donors, a blood hemoglobin level which shall be demonstrated to be no less than 12.5 grams (g) of hemoglobin per 100 milliliters (mL) of blood; or a hematocrit value of 38 percent, and for autologous donors, a blood hemoglobin level which shall be demonstrated to be no less than 11.0 g of hemoglobin per 100 mL of blood or a hematocrit value of 33 percent.

(4) Freedom from acute respiratory diseases;

(5) Freedom from any infectious skin disease at the site of phlebotomy and from any such disease generalized to such an extent as to create a risk of contamination of the blood;

(6) Freedom from any disease transmissible by blood transfusion, insofar as can be determined by history and examinations indicated above; and

(7) Freedom of the arms and forearms from skin punctures or scars indicative of addiction to self-injected narcotics.

(c) *Additional qualifications of donor; viral hepatitis.* No individual shall be used as a source of Whole Blood if he has—

(1) A history of viral hepatitis after the 11th birthday;

(2) A history of close contact within 12 months of donation with an individual having viral hepatitis;

(3) A history of having received within 12 months of donation, human blood or any derivative of human blood which the Food and Drug Administration has advised the blood establishment is a possible source of viral hepatitis.

(d) *Therapeutic bleedings.* Blood withdrawn in order to promote the health of a donor otherwise qualified under the provisions of this section, shall not be used as a source of Whole Blood unless the container label conspicuously indicates the donor's disease that necessitated withdrawal of blood.

(e) [Reserved]

(f) *Qualifications; donations within less than 8 weeks.* A person may serve as a source of Whole Blood more than once in 8 weeks only if at the time of donation the person is examined and certified by a physician to be in good

health, as indicated in part in paragraph (b) of this section.

[38 FR 32089, Nov. 20, 1973, as amended at 49 FR 23834, June 8, 1984; 50 FR 4138, Jan. 29, 1985; 51 FR 15611, Apr. 25, 1986; 55 FR 11013, Mar. 26, 1990; 64 FR 45371, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001; 66 FR 40889, Aug. 6, 2001]

§ 640.4 Collection of the blood.

(a) *Supervision.* Blood shall be drawn from the donor by a qualified physician or under his supervision by assistants trained in the procedure. A physician shall be present on the premises when blood is being collected, except that blood may be collected when a physician is not present on the premises, provided the establishment (1) maintains on the premises, and files with the Center for Biologics Evaluation and Research, a manual of standard procedures and methods, approved by the Director of the Center for Biologics Evaluation and Research, that shall be followed by employees who collect blood, and (2) maintains records indicating the name and qualifications of the person immediately in charge of the employees who collect blood when a physician is not present on the premises.

(b) *The donor center.* The pertinent requirements of §§ 600.10 and 600.11 of this chapter shall apply at both the blood establishment and at any other place where the bleeding is performed.

(c) *Blood containers.* Blood containers and donor sets shall be pyrogen-free, sterile and identified by lot number. The amount of anticoagulant required for the quantity of blood to be collected shall be in the blood container when it is sterilized. In addition, all container and donor set surfaces that come in contact with blood used in the processing of Heparin Whole Blood shall be water repellent.

(d) *The anticoagulant solution.* The anticoagulant solution shall be sterile and pyrogen-free. Anticoagulant solutions shall be compounded and used according to a formula approved by the Director, Center for Biologics Evaluation and Research.

(e) *Donor identification.* Each unit of blood shall be so marked or identified by number or other symbol as to relate it to the individual donor whose iden-

tity shall be established to the extent necessary for compliance with § 640.3.

(f) *Prevention of contamination of the blood.* The skin of the donor at the site of phlebotomy shall be prepared thoroughly and carefully by a method that gives maximum assurance of a sterile container of blood. The blood shall be collected by aseptic methods in a sterile system which may be closed or may be vented if the vent protects the blood against contamination.

(g) *Samples and segments for laboratory tests.* Samples and segments for laboratory tests shall meet the following standards:

(1) One or more segments shall be provided with each unit of blood when issued or reissued except as provided in § 640.2(c)(2) and all segments shall be from the donor who is the source of the unit of blood.

(2) All samples for laboratory tests performed by the manufacturer and all segments accompanying a unit of blood shall be collected at the time of filling the original blood container.

(3) All containers for all samples shall bear the donor's identification before collecting the samples.

(4) All segments accompanying a unit of blood shall be attached to the whole blood container before blood collection, in a tamperproof manner that will conspicuously indicate removal and reattachment.

(5) Segments for compatibility testing shall contain blood mixed with the appropriate anticoagulant.

(h) *Storage.* Whole Blood must be placed in storage at a temperature between 1 and 6 °C immediately after collection unless the blood is to be further processed into another component or the blood must be transported from the donor center to the processing laboratory. If transported, the blood must be placed in temporary storage having sufficient refrigeration capacity to cool the blood continuously toward a temperature range between 1 and 10 °C until arrival at the processing laboratory. At the processing laboratory, the blood must be stored at a temperature between 1 and 6 °C. Blood from which a component is to be prepared must be held in an environment maintained at a temperature range specified for that component in the directions for use for